

STAT 88: Lecture 15

Contents

Section 5.4: Unbiased Estimators

Section 5.5: Conditional Expectation

Section 5.6: Expectation by Conditioning

Last time

Unbiased estimators

Probability distributions often have **parameters** that we wish to estimate. An **estimator** is a **random variable** and there is uncertainty what you will get. With an **unbiased estimator**, on average the estimator will be correct.

Ex Suppose the population has population mean μ , i.e. any sample X from the population has mean $E(X) = \mu$. Let X_1, \dots, X_n be a SRS from the population. We use the sample mean \bar{X} as an estimator of the population mean μ . Sample mean is **always unbiased** since

$$E(\bar{X}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n} \sum_{i=1}^n E(X_i) = \mu.$$

Ex Suppose the **population consists of zeros and ones**. Then the population mean p is the population proportion of ones. If X_1, \dots, X_n are i.i.d. samples from the population, the sample mean \bar{X} is the sample proportion of ones in your sample. By unbiasedness of the sample mean, we have


$$E(\bar{X}) = p.$$

The sampling distribution of sample proportions from the 20,000 repeated experiments:

$n = 30$

$p = 0.1667$

Average of observed sample proportions = 0.1664



$p = \text{population proportion}$
 $= E(\bar{x})$

Warm up: (Exercise 5.7.11) Let X be the number of cars owned by a Cal student. Here is the distribution of X .

number of cars	0	1	2
probability	2θ	θ	$1 - 3\theta$

(a) Find $E(X)$ (as a function of θ).

(b) Let X_1, \dots, X_n be the number of cars owned by n randomly picked students. Use \bar{X} to find an unbiased estimator of θ .

5.4. Unbiased Estimators (Continued)

Estimating the largest possible value

Let $X_1, X_2, \dots, X_n \stackrel{i.i.d.}{\sim} \text{Uniform}\{1, 2, \dots, N\}$ for some fixed but unknown N . To estimate N , there are two possible estimators we can come up with:


parameter

1. $M = \max\{X_1, \dots, X_n\}$. Note that this is a biased estimator.
2. We know that the population mean is $\mu = (N+1)/2$ and thus $E(\bar{X}) = (N+1)/2$ since it is unbiased. Then what is an estimator T such that

$$E(T) = N?$$

Lets look at sampling distribution of (1) $M = \max(X_1, \dots, X_n)$ and (2) $T = 2\bar{X} - 1$.

N = 300
n = 30
5000 repetitions



The histograms show that both estimators have pros and cons.

M - Pros: small spread of values; Cons: biased.

T - Pros: unbiased; Cons: big spread of values.

Unbiasedness is a good property, but so is low variability. Bias-variance tradeoff

5.5. Conditional Expectation

Let's first review how to find expectation of a joint distribution. A joint distribution for two random variables X and S is given below:

	$X = 1$	$X = 2$	$X = 3$
$S = 2$	0.0625	0	0
$S = 3$	0.125	0.125	0
$S = 4$	0.0625	0.25	0.0625
$S = 5$	0	0.125	0.125
$S = 6$	0	0	0.0625

The marginal distribution of S is given by summing along the rows:

s	2	3	4	5	6
$P(S = s)$	0.0625	0.25	0.375	0.25	0.0625

Conditional Distribution Suppose someone runs the experiment and tells you that $S = 3$. Given this information, what is the distribution of X ?

$$P(X = 1|S = 3) = \frac{P(X = 1, S = 3)}{P(S = 3)} = \frac{0.125}{0.25} = 0.5.$$

Similarly we can get $P(X = 2|S = 3) = 0.5$ and $P(X = 3|S = 3) = 0$.

If X and S are two random variables on the same outcome space, then for a fixed value s of S , the conditional distribution of X given $S = s$ is

- the set of all possible values of X under the condition that $S = s$, and
- all the corresponding conditional probabilities $P(X = x|S = s)$.

The distribution of X changes depending on the given value of S :

	$X=1$	$X=2$	$X=3$		$X=1$	$X=2$	$X=3$
Conditional Dist'n of X given $S=3$	0.5	0.5	0	Conditional Dist'n of X given $S=4$	0.667	0.667	0.667

Conditional Expectation The expectation of X , also called the unconditional expectation of X , is easy to see from the distribution table:

x	1	2	3
$P(X = x)$	0.25	0.5	0.25

$$E(X) =$$

Given that S has the value s , the conditional distribution of X is just an ordinary distribution and thus has an expectation. This is called the conditional expectation of X given $S = s$ and is denoted $E(X|S = s)$.

$$E(X|S = 3) =$$

Generally, $E(X|S=s) = \sum_{\text{all } x} x P(X=x|S=s)$

Unconditional Expectation
 $E(X) = \sum_{\text{all } x} x P(X=x)$

What is relationship between expectation and conditional expectation?

$$E(X) = \sum_{\text{all } x} xP(X = x) = \sum_{\text{all } x} \sum_{\text{all } s} xP(X = x, S = s).$$

By multiplication rule,

$$P(X = x, S = s) = P(X = x|S = s)P(S = s).$$

So

$$E(X) = \sum_{\text{all } s} \sum_{\text{all } x} xP(X = x, S = s) = \sum_{\text{all } s} \underbrace{\sum_{\text{all } x} xP(X = x|S = s)}_{=E(X|S=s)} P(S = s).$$

Therefore

$$E(X) = \sum_{\text{all } s} E(X|S = s)P(S = s).$$

Important: $E(X|S = s)$ is a function of s . For example,

s	2	3	4	5	6
$E(X S = s)$	1	1.5	2	2.5	3

Marginal Distribution of S

s	2	3	4	5	6
$P(S = s)$	0.0625	0.25	0.375	0.25	0.0625

5.6. Expectation by Conditioning

To find expectation of one random variable, it sometimes helps to condition on another random variable.

Time to Reach Campus A student has two routes to campus. Each route has a random duration. The student prefers Route A because its expected duration is 15 minutes compared to 20 minutes for Route B. However, on 10% of her trips she is forced to take Route B because of road work on Route A. On the remaining 90% of the days she takes Route A.

What is the expected duration of her trip on a random day?

Example: (Exercise 5.7.13) A survey organization in a town is studying families with children. Among the families that have children, the distribution of the number of children is as follows.

Number of Children n	1	2	3	4	5
Proportion with n Children	0.2	0.4	0.2	0.15	0.05

Suppose each child has chance 0.51 of being male, independently of all other children. What is the expected number of male children in a family picked at random from those that have children?

Example: You flip a fair coin N times where N is a random variable $N \sim \text{Poisson}(5)$.
What is the expected number of heads you will get?